

Magnet Shuffling for the RHIC

S. Ohnuma

November 12, 1985

Magnet Shuffling for the RHIC (?)

1. Remarks

- a) Correction Systems (What type, How many families, How to tune them, What on Day One).
- b) Magnets in the insertions are special.
special consideration at the colliding points.

skew quadrupoles at Q2 or Q3 to control
the coupling, $v_x - v_y = 0$.

At Q2 or Q3.

- (1) $\beta^* = 3\text{m}$, two families are orthogonal.
- (2) $\beta^* = 6\text{m}$, two families are NOT independent.

$$\underbrace{(\psi_x - \psi_y) - (v_x - v_y) \cdot \theta}_{}$$

2. "Local" vs "Global"

"closed Form" vs "Fourier Series"

- Example
- 1. isolated resonances, resonance extraction.
 - 2. localized source of errors
(e.g. elements in insertions)

① $\Delta \hat{X}_p$ by b_1

$$(\Delta \hat{X}_p) = \frac{\Delta X_p}{\sqrt{\beta_x}} + i < \Delta \hat{X}_p' + \frac{\alpha_x}{\beta_x} \cdot \Delta X_p > \cdot \sqrt{\beta_x}$$

"Global" (b_1 distributed around the ring)


$$(\Delta \hat{X}_p)_\phi = \sqrt{\beta_x(\phi)} \cdot V_x^z \cdot \sum_{-\infty}^{\infty} \frac{a_n}{V_x^z - n^2} \cdot e^{in\phi_x}; \phi = \psi/\nu$$

$$a_n = \frac{1}{2\pi V_x} \cdot \sum_k^N \left[\left(-\frac{\ell}{\rho} \right) \cdot b_1 \cdot X_p \cdot \sqrt{\beta_x} e^{-in\phi_x} \right]_k$$

Error b_1 at $k=1$ to N .

"Local" (b_1 confined to localized area)

OUTSIDE vs INSIDE



$$(\Delta \hat{X}_p)_o = \frac{e^{-i\pi\nu}}{2\sin(\pi\nu)} \cdot \sum_k^M \left[\left(-\frac{\ell}{\rho} \right) b_1 \sqrt{\beta_x} X_p \cdot e^{i\psi_x} \right]_k$$

If $\left(-\frac{\ell}{\rho} \right) \cdot \sqrt{\beta_x} \cdot X_p$ is the same at $k=1 \sim M (\ll N)$

$$(\Delta \hat{X}_p)_o = 0 \rightarrow \boxed{\sum_k^M (b_1 \cdot e^{i\psi_x})_k = 0}$$

Confine the effect within the localized area.

③ γ_p by a_1

$$x \rightarrow y; b_1 \rightarrow a_1$$

$$\text{"Local"} \quad \left[\left(\frac{\ell}{p} \right) \cdot a_1 \cdot X_p \cdot \sqrt{\beta y} \cdot e^{i\psi_y} \right]_k \rightarrow \sum_k^M (a_1 e^{i\psi_y})_k = 0$$

$$\text{"Global"} \quad \left[\left(\frac{\ell}{p} \right) \cdot a_1 \cdot X_p \cdot \sqrt{\beta y} \cdot e^{-in\phi_y} \right]_k$$

③ $(\Delta\beta/\beta)_{x+y}$ by b_1

$$\hat{A} \quad \hat{A} = (\Delta\beta/\beta) - i \langle \Delta\alpha - \frac{\Delta\beta}{\beta} \cdot \alpha \rangle$$

$$\text{"Global"} \quad \hat{A} = (2\nu^2) \cdot \sum_{-\infty}^{\infty} \frac{J_n}{4\nu^2 - n^2} \cdot e^{in\phi}$$

$$J_n = \frac{1}{2\pi\nu} \cdot \sum_k^N \left[\left(-\frac{\ell}{p} \right) \cdot b_1 \cdot \beta \cdot e^{-in\phi} \right]_k$$

"Local"

$$\hat{A}(\text{OUTSIDE}) = \frac{e^{-i2\pi\nu}}{2\sin(2\pi\nu)} \cdot \sum_{k=1}^M \left[\left(-\frac{\ell}{p} \right) \cdot \beta \cdot b_1 \cdot e^{i2\psi} \right]_k$$

$$\hat{A} = 0 \quad \rightarrow \boxed{\sum_k^M (b_1 \cdot e^{i2\psi})_k = 0.}$$

Two ways of looking at the same problem.

(1) "Local".

Confine the effect such that no effect OUTSIDE.

a) Given ~~N~~ N, what M?

b) Combination of two or more blocks?


(2) "Global"

Reduce the "important" harmonics.

($n \approx v$ or $n \approx 2v$).

How many harmonics? \rightarrow Again localized area.

3. Thin-lens approximation




Can we take each dipole as an element or do we have to regard each dipole as several elements?

	thin-lens	integrated
$\sqrt{\beta_x} \cdot X_p \cdot e^{i\psi_x}$:	4.95 (12.5°)	5.15 (11.5°)
$\sqrt{\beta_y} \cdot X_p \cdot e^{i\psi_y}$:	4.95 (33.0°)	4.89 (31.9°)
$\beta_x \cdot e^{iz\psi_x}$:	22.4 (25.0°)	22.8 (22.5°)
$\beta_y \cdot e^{iz\psi_y}$:	22.4 (65.9°)	22.8 (68.4°)


distortion by b_2

$\beta_x^{3/2} e^{i3\psi_x}$	106.0 (37.5°)	112.6 (30.3°)
------------------------------	------------------------	------------------------


Should be OK for (b_1, q_1) but may have to shift the angle for (b_2, q_2) .




4. Example of "Vector Diagrams" for RHIC.



$a_1 \sqrt{p_y} X_p e^{i\psi_y}$





Important differences from the Tevatron

(1) Tevatron : b_2, q_2, q_3

(2) RHIC : $b_1, q_1, b_2, q_2 (?)$

Phase advance / cell. RHIC : 90°

Tevatron : 68°

For RHIC, various effects coming from one multipole (for example, b_1) are correlated.

5. What I should do

Given $\langle b_1 \rangle, \langle q_1 \rangle, \langle b_2 \rangle \& \langle q_2 \rangle$.

- (1) max. allowed values (if Gaussian)
- (2) $M=8$ (~~two~~ four cells) or 16 (eight cells)
- (3) "manual" and "automated" shuffling processes combined.
- (4) performance estimates.

Compare with random cases:

$(\Delta x_p)_{\text{max.}}$	$(Y_p)_{\text{max.}}$	$(\beta^2/\beta)_{\text{max.}}$
$\langle \Delta x_p \rangle_{\text{RMS.}}$	"	"

Distortion functions - - -